

On chromatic number of unit-quadrance graphs (finite Euclidean graphs)

Le Anh Vinh
 School of Mathematics
 University of New South Wales
 Sydney 2052 NSW

Abstract

The quadrance between two points $A_1 = (x_1, y_1)$ and $A_2 = (x_2, y_2)$ is the number $Q(A_1, A_2) = (x_1 - x_2)^2 + (y_1 - y_2)^2$. Let q be an odd prime power and F_q be the finite field with q elements. The unit-quadrance graph D_q has the vertex set F_q^2 , and $X, Y \in F_q^2$ are adjacent if and only if $Q(A_1, A_2) = 1$. Let $\chi(F_q^2)$ be the chromatic number of graph D_q . In this note, we will show that $q^{1/2}(1/2 + o(1)) \leq \chi(F_q^2) \leq q(1/2 + o(1))$. As a corollary, we have a construction of triangle-free graphs D_q of order q^2 with $\chi(D_q) \geq q/2$ for infinitely many values of q .

1 Introduction

In [3], Wildberger introduces a remarkable new approach to trigonometry and Euclidean geometry by replace distance by quadrance and angle by spread, thus allowing the development of Euclidean geometry over any field. The following definition follows from [3].

Definition 1 *The **quadrance** $Q(A_1, A_2)$ between the points $A_1 = (x_1, y_1)$, and $A_2 = (x_2, y_2)$ in F_q^2 is the number*

$$Q(A_1, A_2) := (x_2 - x_1)^2 + (y_2 - y_1)^2.$$

This approach motives the question of finding chromatic number of unit-quadrance graph over any field. Note that in usual 2-dimensional Euclidean space R^2 then the quadrance between A_1, A_2 is unit if and only if the distance between A_1, A_2 is unit.

Let q be an odd prime power and F_q be the finite field with q elements. The unit-quadrance graph D_q has the vertex set F_q^2 , and $X, Y \in F_q^2$ are adjacent if and only if $Q(A_1, A_2) = 1$. This graph (or so-called finite Euclidean graph) was also studied by Medrano et al in [2]. Recall that the graph D_q is a regular graph with degree $\Delta(D_q) = q - (-1)^{(q-1)/2}$ (see [2]). Let $\chi(F_q^2)$ be the chromatic number of graph D_q . The main result of this note is the following theorem.

Theorem 1 Suppose that $q = p^n > 3$ where p is an odd prime number then

$$q^{1/2}(1/2 + o(1)) \leq \chi(F_q^2) \leq \frac{p^n + p^{n-1}}{2} = q(1/2 + o(1)).$$

2 Some Lemmas

In order to prove Theorem 1, we need some lemmas.

Lemma 1 Suppose that $a \in F_q$ such that $a^2 + 1$ is not square in F_q . For any $A \neq B$ in the line $y = ax + i$ then $Q(A, B) \neq 1$.

Proof Suppose that $A = (x_1, ax_1 + i)$ and $B = (x_2, ax_2 + i)$ for some $x_1 \neq x_2 \in F_q$. We have

$$Q(A, B) = (a^2 + 1)(x_1 - x_2)^2 \neq 1$$

since $a^2 + 1$ is not square in F_q . The lemma follows. $\square$

Recall that a (multiplicative) character of F_q is a homomorphism from F_q^* , the multiplicative group of the non-zero elements of F_q , to the multiplicative group of complex numbers with modulus 1. The identically 1 function is the principal character of F_q and is denoted χ_0 . Since $x^{q-1} = 1$ for every $x \in F_q^*$ we have $\chi^{q-1} = \chi_0$ for every character χ . A character χ is of order d if $\chi^d = \chi_0$ and d is the smallest positive integer with this property. By convention, we extend a character χ to the whole of F_q by putting $\chi(0) = 0$. The quadratic (residue) character is defined by $\chi(x) = x^{(q-1)/2}$. Equivalently, χ is 1 on square, 0 at 0 and -1 otherwise. It is easy to see that $\sum_{i \in F_q} \chi(i) = 0$. We have the following lemma.

Lemma 2 Suppose that t is not a square in F_q . Then there exists a square number $i \in F_q^*$ such that $-i + t$ is not a square in F_q .

Proof Let A_q be the number of $i \in F_q^*$ such that i is square but $-i + t$ is not. Consider the sum

$$\begin{aligned} \sum_{i \in F_q} (1 - \chi(i)) (1 + \chi(-i + t)) &= q + \sum_{i \in F_q} \{-\chi(i) + \chi(-i + t) - \chi(i(-i + t))\} \\ &= q - \sum_{i \in F_q \neq 0, t} \chi(i^2(-1 + ti^{-1})) \\ &= q - \sum_{i \in F_q \neq 0, t} \chi(-1 + ti^{-1}) \\ &= q + \chi(-1) + \chi(0) \\ &= q + (-1)^{(q-1)/2}. \end{aligned}$$

But we have

$$(1 - \chi(i))(1 + \chi(-i + t)) = \begin{cases} 4 & \text{if } i \text{ is square and } -i + t \text{ is not square,} \\ 2 & \text{if } i = t. \\ 0 & \text{otherwise.} \end{cases}$$

Thus, $4A_q + 2 = q + (-1)^{(q-1)/2}$ or $A_q = (q + (-1)^{(q-1)/2} - 2)/4 \geq 1$. This concludes the proof of the lemma. $\square$

Lemma 3 *Suppose that $q \geq 5$ is a prime power. Let $a \in F_q$ such that $a^2 + 1$ is not square. There exist $t \in F_q^*$ such that if A is in the line $y = ax + i$ and B is in the line $y = ax + i + t$ then $Q(A, B) \neq 1$ for any $i \in F_q$.*

Proof Suppose that $X = (x, ax + i)$ and $Y = (y, ay + i + t)$ then

$$\begin{aligned} Q(X, Y) &= (x - y)^2 + (a(x - y) + t)^2 \\ &= (a^2 + 1)(x - y)^2 + 2a(x - y)t + t^2. \end{aligned}$$

If $Q(X, Y) = 1$ then $a^2 + 1 = [(a^2 + 1)(x - y) + at]^2 + t^2$. From Lemma 2, there exists $i \in F_q^*$ such that i is square but $-i + a^2 + 1$ is not. Hence, we can choose t such that $t^2 = i$. This concludes the proof of the theorem. $\square$

In [2], Medrano et al. give a general bound for eigenvalue of D_q .

Lemma 4 ([2]) *Let $\lambda \neq \Delta(D_q)$ be any eigenvalue of graph D_q then $|\lambda| \leq q^{1/2}$.*

The following result which is due to Hoffman gives us a connection between the chromatic number and eigenvalues of a graph.

Lemma 5 ([1]) *Let G be any graph with the largest and least eigenvalues are λ_1 and λ_* . Then*

$$\chi(G) \geq 1 - \frac{\lambda_1}{\lambda_*}.$$

3 Proof of Theorem 1

The lower bound is straightforward from Lemmas 4 and 5. We have $\lambda_1 = q \pm 1$ since D_q is a $(q \pm 1)$ -regular graph. Lemma 4 gives us $\lambda_* \geq -q^{1/2}$. Hence

$$\chi(D_q) \geq 1 + \frac{q \pm 1}{q^{1/2}} = q^{1/2}(1 + o(1)).$$

The upper bound is proved by a geometrical colouring. Let $a \in F_q^*$ such that $a^2 + 1$ is not a square in F_q and choose t satisfy Lemma 3. We partition F_q into p^{n-1} disjoint sets of the form $\{\alpha_i, \alpha_i + t, \dots, \alpha_i + (p-1)t\}$ for $i = 1, \dots, p^{n-1}$. We colour $2p$ points in the lines

3	1	2	3	4	1	2
2	3	4	1	2	3	1
4	1	2	3	1	2	3
2	3	1	2	3	4	1
1	2	3	4	1	2	3
3	4	1	2	3	1	2
1	2	3	1	2	3	4

Table 1: 4-colouring of F_7^2

$y = ax + \alpha_i + (2k)t$ and $y = ax + \alpha_i + (2k+1)t$ the same colour for each $i = 1, \dots, p^{n-1}$ and $0 \leq k \leq (p-3)/2$ (using different colours for each pair (i, k)). And we colour p points in the line $y = ax + \alpha_i + p - 1$ the same colour for each $i = 1, \dots, p^{n-1}$ (using different colour for each i). Then total number of colours we have used is $p^{n-1}(p+1)/2$. From Lemmas 1 and 3, there does not exist two point A, B such that $Q(A, B) = 1$ and A, B have the same colour. This implies the upper bound and concludes the proof of the theorem.

Example 1 For $q = 7$ then we have $3 \leq \chi(F_7^2) \leq 4$ (the upper bound follows from Theorem 1, the lower bound is trivial since D_7 contains a cycle of length 7). We choose $a = 5, t = 3$ then $a^2 + 1 = 5$ and $-t^2 + a^2 + 1 = 3$ are not square in F_7 . We have a 4-colouring of points of F_7^2 in Table 1. It has been verified by computer that there does not exist a 3-colouring of vertices of D_7 . This implies that $\chi(F_7^2) = \chi(D_7) = 4$.

4 Triangle-free graphs with arbitrary high chromatic numbers

The unit-quadrance graphs give us a family of triangle-free graphs with arbitrary high chromatic numbers. We have the following lemma.

Lemma 6 Let q be any prime of the form $q = 12k \pm 7$. Then D_q contains no triangle.

Proof Suppose that D_q contains a triangle XYZ with $X = (m, n)$, $Y = (m+x, n+y)$ and $Z = (m+x+u, n+y+v)$ for some $m, n, x, y, u, v \in F_q$ then $x^2 + y^2 = u^2 + v^2 = (x+u)^2 + (y+v)^2 = 1$. This implies that $xu + yv = -1/2$. We have $(xu+yv)^2 + (xv-yu)^2 = (x^2 + y^2)(u^2 + v^2)$ so

$$(xv - yu)^2 = 1 - 1/4 = 3/4.$$

For $q = 12 \pm 7$ then 3 is not square in F_q . Hence we have a contradiction. This concludes the proof of the lemma. $\square$

From Theorem 1 and Lemma 6, if $q \equiv \pm 7 \pmod{12}$ is a prime number then D_q is a triangle-free graph with chromatic number $\chi(D_q) \geq q/2(1 + o(1))$. This give us a family of triangle-free graphs with arbitrary high chromatic numbers.

5 Higher dimensional spaces

The quadrance can be defined in higher dimension space F_q^m for $m \geq 2$ as follows.

Definition 2 The **quadrance** $Q(A_1, A_2)$ between the points $X = (x_1, \dots, x_m)$, $Y = (y_1, \dots, y_m) \in F_q^m$ is the number

$$Q(A_1, A_2) := \sum_{i=1}^m (x_i - y_i)^2.$$

The unit-quadrance graph D_q^m has the vertex set F_q^m , and $X, Y \in F_q^m$ are adjacent if and only if $Q(A_1, A_2) = 1$. Let $\chi(F_q^m)$ be the chromatic number of graph D_q^m . Similar as the above, we have the following theorem.

Theorem 2 Suppose that $m \geq 2$ and $q = p^n > 3$ where p is an odd prime number then

$$q^{(m-1)/2}(1/2 + o(1)) \leq \chi(F_q^m) \leq \frac{q^{m-2}[p^n + p^{n-1}]}{2} = q^{m-1}(1/2 + o(1)).$$

The proof of this theorem is omitted since it is the same as the proof of Theorem 1.

References

- [1] Hoffman, A. J., On eigenvalues and colorings of graphs, in *Graph Theory and Its Applications* (B. Harris, ed.), Acad. Press, 1970, 79-91.
- [2] Medrano, A., Myers, P., Stark, H. M., Terras, A., Finite analogues of Euclidean space, *Journal of Computational and Applied Mathematics*, **68** (1996), 221-238
- [3] Wildberger, N. J., *Divine Proportions: Rational trigonometry to universal geometry*, WildEgg, 2005.